

# King's College London

UNIVERSITY OF LONDON

**This paper is part of an examination of the College counting towards the award of a degree. Examinations are governed by the College Regulations under the Authority of the Academic Board.**

**M.Sci. EXAMINATION**

**CP/4730 The C programming language for physicists**

**SUMMER 1998**

Time allowed: **TWO HOURS**

**Candidates must answer any TWO questions. No credit will be given for attempting a further question.**

**The approximate mark for each part of a question is indicated in square brackets.**

**Good answers to questions will include plans and explanations in addition to sections of C code.**

**TURN OVER WHEN INSTRUCTED**

© 1998 King's College London

## Answer TWO questions

- 1) Write a short program in C, which reads in an integer and finds all of its prime factors, including repeated factors.

[20 marks]

- 2) The diffusion equation in one dimension is given by:

$$D \frac{\partial^2 C}{\partial x^2} = \frac{\partial C}{\partial t}$$

where  $D$  is the diffusion constant of the material and  $C$  is the concentration of the diffusing quantity.

Write a short C program which reads an array from a file `initial.d`, which contains the concentration  $C(x,0)$  at time zero, at  $n$  equally and closely spaced points along the  $x$ -axis, and calculates the concentration,  $C(x,t)$ , at later times using the finite difference expressions for the differentials in the diffusion equation. Set the values at the ends of the  $x$ -axis to be constant in time.

[20 marks]

[For a discrete time step  $dt$ :  $\frac{\partial C}{\partial t} \approx \frac{C(x,t+dt) - C(x,t)}{dt}$ , but for the spatial

differentials, use the symmetrical version:

$$\frac{\partial^2 C}{\partial x^2} \approx \frac{C(x+dx,t) + C(x-dx,t) - 2C(x,t)}{dx^2}$$

where  $dx$  is the distance between the points at which  $C$  is defined.]

- 3) Write a function in C which calculates the function  $\text{sinc}(x) = \frac{\sin x}{x}$  to 5 significant figures, using the series for  $\sin x$ :

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \mathbf{L}$$

You should use some sort of convergence criterion, and be especially careful when  $|x|$  is large or very near zero.

[20 marks]

**FINAL PAGE**